SYMMETRY REDUCTIONS AND EXACT SOLUTIONS OF LAX INTEGRABLE 3-DIMENSIONAL SYSTEMS

H. BARAN, I.S. KRASIL′SHCHIK, O.I. MOROZOV, AND P. VOJČÁK

Abstract. We present a complete description of 2-dimensional equations that arise as symmetry reductions of fourf 3-dimensional Lax-integrable equations: (1) the universal hierarchy equation $u_{yy} = u_z u_{xy} - u_y u_{xz}$; (2) the 3D rdDym equation $u_{ty} = u_x u_{xy} - u_y u_{xx}$; (3) The basic Veronese web equation $u_{ty} = u_t u_{xy} - u_y u_{tx}$; (4) Pavlov's equation $u_{yy} = u_{tx} + u_y u_{xx} - u_x u_{xy}$.

Contents

Introduction 2
1. Preliminaries 3
2. The universal hierarchy equation 3
2.1. Symmetries 3
2.2. Reductions 4
3. The 3D rdDym equation 7
3.1. Symmetries 7
3.2. Reductions 7
4. The basic Veronese web equation 11
4.1. Symmetries 11
4.2. Reductions 12
5. Pavlov's equation 14
5.1. Symmetries 14
5.2. Reductions 15
6. Summary of results 18
Acknowledgements 18
References 19

arXiv:1407.0246v1 [nlin.SI] 1 Jul 2014

Date: July 2, 2014.
2010 *Mathematics Subject Classification.* 35B06.
Key words and phrases. Partial differential equations, symmetry reductions, solutions.

Introduction

We consider the four 3-dimensional Lax-integrable[1] equations:

- the *universal hierarchy equation* (Section 2)
$$u_{yy} = u_z u_{xy} - u_y u_{xz}, \tag{1}$$
see [11].
- the *3D rdDym equation* (Section 3)
$$u_{ty} = u_x u_{xy} - u_y u_{xx}, \tag{2}$$
see [2, 13, 16].
- the *basic Veronese web equation* (Section 4)
$$u_{ty} = u_t u_{xy} - u_y u_{tx}, \tag{3}$$
see [1, 5, 7, 14].
- *Pavlov's equation* (Section 5)
$$u_{yy} = u_{tx} + u_y u_{xx} - u_x u_{xy}, \tag{4}$$
see [4, 15].

Some of these equations arise also in [6] as integrable hydrodynamic reductions of multi-dimensional dispersionless PDEs.

All the above listed equations may be obtained as the symmetry reductions of the following Lax-integrable 4-dimensional systems:

$$u_{yz} = u_{tx} + u_x u_{xy} - u_y u_{xx}$$

and

$$u_{ty} = u_z u_{xy} - u_y u_{xz}$$

introduced in [6] and [11], respectively, while the latter two, in turn, are the reductions of

$$u_{yz} = u_{ts} + u_s u_{xz} - u_z u_{xs}.$$

Here we give a complete answer to a natural question: what 2-dimensional equations are the reductions of the 3-dimensional ones? The result comprises 32 equations of which

- sixteen can be solved explicitly,
- one reduces to the Riccati equation,
- five can be linearized by the Legendre transformation,
- while the rest ten are 'nontrivial'.

The latter are presented in Table 1 (in the third column, we exemplify the simplest relations). The first two of these equations can be transformed to the Liouville equation and the Gibbons-Tsarev equation, respectively. The other eight, to our strong opinion, may possess interesting integrability properties and we plan to study them in the nearest future. More detailed, but concise, infomation on the reductions may be also found in

Reduction	of Eq.	Relations with the initial equation
$2\Phi = \Phi\Phi_{xz} - \Phi_x\Phi_z,$	(1)	$u = \dfrac{\Phi(x,z)}{y},$
$\Phi_{\xi\xi} = (\xi + \Phi_\xi)\Phi_{\eta\eta} - \Phi_\eta\Phi_{\xi\eta} - 2,$	(4)	$u = \Phi(\xi,\eta) + t^2\xi - 2t\eta,\ \xi = y,\ \eta = x + ty,$
$\Phi_{\xi\xi} = \Phi_x\Phi_\xi - \Phi\Phi_{x\xi},$	(1)	$u = \Phi(x,\xi)e^{-z},\ \xi = ye^{-z},$
$(1 + \xi\Phi_z)\Phi_{\xi\xi} - \xi\Phi_\xi\Phi_{\xi z} + \Phi_\xi\Phi_z = 0,$	(1)	$u = \Phi(z,\xi)e^{-x},\ \xi = ye^{-x},$
$\Phi_\eta\Phi_{\xi\eta} - \Phi_\xi\Phi_{\eta\eta} = e^\eta\Phi_{\xi\xi},$	(1)	$u = \Phi(\xi,\eta)e^{-x},\ \xi = ye^{-z},\ \eta = x - z,$
$(\xi + \Phi_\xi)\Phi_{\xi y} - \Phi_y(\Phi_{\xi\xi} + 2) = 0,$	(2)	$u = \Phi(\xi,y)e^{2t},\ \xi = xe^t,$
$\Phi_{\xi t} = 4\Phi\Phi_\xi - \xi\Phi_\xi^2 + 2\xi\Phi\Phi_{\xi\xi},$	(2)	$u = \Phi(\xi,t)x^2,\ \xi = xe^{-y},$
$\Phi_{\eta\eta} + (\xi + \Phi_\eta)\Phi_{\xi\eta} = \Phi_\eta(2 + \Phi_{\xi\xi}),$	(2)	$u = \Phi(\xi,\eta)e^{2t},\ \xi = xe^{-t},\ \eta = y - t,$
$(4\xi^2 - 3\Phi)\Phi_{\xi\xi} - \Phi_{\xi t} - 6\xi\Phi_\xi + \Phi_\xi^2 + 6\Phi = 0,$	(4)	$u = \Phi(\xi,y)y^3,\ \xi = \dfrac{x}{y^2},$
$\Phi_{\xi\xi} = (\xi - \Phi_\eta)\Phi_{\xi\eta} + (2\eta + \Phi_\xi)\Phi_{\eta\eta} - \Phi_\eta = 0,$	(4)	$u = \Phi(\xi,\eta)e^{-3t},\ \xi = ye^{\beta t}, \eta = xe^{2t}$

Table 1. 'Nontrivial' reductions

[1]We say that an equation is *Lax-integrable* if it admits a zero-curvature representation with a non-removable parameter.

Table 6.

In Section 1, we briefly expose necessary preliminaries (see, e.g., [10]). In Section 6, we present the obtained results in a concise form.

1. Preliminaries

Let $\mathscr{E}$ be a differential equation given by

$$F\Big(x,\dots,\frac{\partial^{|\sigma|}u}{\partial x^\sigma},\dots\Big)=0, \tag{5}$$

where $u(x)$ is the unknown function in the variables $x=(x^1,\dots,x^n)$. A *symmetry* of $\mathscr{E}$ is a function $\varphi=\varphi(x,\dots,u_\sigma,\dots)$ in the *jet variables* u_σ, σ being a multi-index, $u_\varnothing=u$, that satisfies the *linearized equation*

$$\ell_{\mathscr{E}}(\varphi)\equiv\sum_\sigma\frac{\partial F}{\partial u_\sigma}D_\sigma(\varphi)=0, \tag{6}$$

where $D_\sigma=D_{i_1}\circ\dots\circ D_{i_k}$ for $\sigma=i_1\dots i_k$, while

$$D_i=\frac{\partial}{\partial x^i}+\sum_\sigma u_{\sigma i}\frac{\partial}{\partial u_\sigma} \tag{7}$$

are the *total derivatives* restricted to $\mathscr{E}$. Symmetries of $\mathscr{E}$ form a Lie algebra $\operatorname{sym}\mathscr{E}$ over $\mathbb{R}$ with respect to the *Jacobi bracket*

$$\{\varphi,\bar\varphi\}=\sum_\sigma\left(\frac{\partial\varphi}{\partial u_\sigma}D_\sigma(\bar\varphi)-\frac{\partial\bar\varphi}{\partial u_\sigma}D_\sigma(\varphi)\right). \tag{8}$$

A solution u to Equation (5) is said to be *invariant* with respect to a symmetry $\varphi\in\operatorname{sym}\mathscr{E}$ if it enjoys the equation

$$\varphi\Big(x,\dots,\frac{\partial^{|\sigma|}u}{\partial x^\sigma},\dots\Big)=0. \tag{9}$$

The *reduction* of $\mathscr{E}$ with respect to φ is Equation (5) rewritten in terms of first integrals of Equation (9).

2. The universal hierarchy equation

The equation is

$$u_{yy}=u_zu_{xy}-u_yu_{xz}. \tag{10}$$

2.1. Symmetries.

The defining equation for symmetries for (10) is

$$D_y^2(\varphi)=u_zD_xD_y(\varphi)-u_yD_xD_z(\varphi)+u_{xy}D_z(\varphi)-u_{xz}D_y(\varphi). \tag{11}$$

Its solutions are

$$\begin{aligned}\varphi_1&=yu_y+u,\\ \varphi_2(X_2)&=X_2u_x-X_2'u,\\ \varphi_3(Z_3)&=Z_3u_z+Z_3'yu_y,\\ \varphi_4(Z_4)&=Z_4u_y,\\ \varphi_5(X_5)&=X_5,\end{aligned}$$

where X_i are functions of x, Z_i are functions of z and 'prime' denotes the derivative with respect to the corresponding variable. The commutator relations are given in Table 2

	φ_1	$\varphi_2(\bar X_2)$	$\varphi_3(\bar Z_3)$	$\varphi_4(\bar Z_4)$	$\varphi_5(\bar X_5)$
φ_1	0	0	0	$\varphi_4(\bar Z_4)$	$-\varphi_5(\bar X_5)$
$\varphi_2(X_2)$	…	$\varphi_2(\bar X_2X_2'-X_2\bar X_2')$	0	0	$\varphi_5(\bar X_5X_2'-X_2\bar X_5')$
$\varphi_3(Z_3)$	…	…	$\varphi_3(\bar Y_3Y_3'-Y_3\bar Y_3')$	$\varphi_4(\bar Y_4Y_3'-Y_3\bar Y_4')$	0
$\varphi_4(Z_4)$	…	…	0	0	0
$\varphi_5(X_5)$	…	…	…	…	0

Table 2. Lie algebra structure of $\operatorname{sym}\mathscr{E}_{(10)}$

2.2. **Reductions.** Thus, the general symmetry of Equation (10) is

$$\varphi = X_2 u_x + (\alpha y + Z_3' y + Z_4) u_y + Z_3 u_z + (\alpha - X_2') u + X_5,$$

where $\alpha \in \mathbb{R}$ is a constant. Thus, invariant with respect to φ solutions are given by the system

$$\frac{dx}{X_2} = \frac{dy}{(\alpha + Z_3')y + Z_4} = \frac{dz}{Z_3} = -\frac{du}{(\alpha - X_2')u + X_5}. \tag{12}$$

We consider the following basic cases below:

Case 00**:** $X_2 = 0$, $Z_3 = 0$;
Case 01**:** $X_2 = 0$, $Z_3 \neq 0$;
Case 10**:** $X_2 \neq 0$, $Z_3 = 0$;
Case 11**:** $X_2 \neq 0$, $Z_3 \neq 0$.

2.2.1. *Case* 00. System (12) takes the form

$$\frac{dx}{0} = \frac{dy}{\alpha y + Z_4} = \frac{dz}{0} = -\frac{du}{\alpha u + X_5}.$$

Its integrals are

$$(\alpha y + Z_4)u + X_5 y = \text{const}, \quad x = \text{const}, \quad z = \text{const}$$

and the general solution is given by

$$\Psi((\alpha y + Z)u + Xy, x, z) = 0,$$

where $Z = Z_4$, $X = X_5$. Hence,

$$u = \frac{\Phi(x, z) - Xy}{\alpha y + Z}. \tag{13}$$

To simplify the subsequent exposition, we consider two subcases:

Subcase 00.0**:** $\alpha = 0$;
Subcase 00.1**:** $\alpha \neq 0$.

Subcase 00.0. After redenoting $\Phi \mapsto \Phi/Z$, $Z \neq 0$, we have

$$u = \Phi(x, z) - \frac{Xy}{Z}. \tag{14}$$

Substituting to Equation (10), one obtains

$$\frac{1}{Z} \cdot (X\Phi_{xz} - X'\Phi_z) = 0,$$

which leads to the following class of solutions

$$\boxed{u = \begin{cases} \Phi(x, z), & \text{if } X = 0, \\ XP(z) + Q(x) - \frac{Xy}{Z}, & \text{if } X \neq 0. \end{cases}}$$

Subcase 00.1. Making the change $\Phi \mapsto \Phi - XZ$, one gets

$$u = \frac{\Phi}{y + Z} - X.$$

Substituting to (10), one arrives to the equation

$$\boxed{2\Phi = \Phi\Phi_{xz} - \Phi_x\Phi_z.}$$

After the change $\Phi = e^{\Psi}$ we obtain the Liouville equation

$$\boxed{\Psi_{xz} = 2e^{-\Psi},}$$

see, e.g. [3].

2.2.2. *Case* 01. Now we have

$$\frac{dx}{0}=\frac{dy}{(\alpha+Z_3')y+Z_4}=\frac{dz}{Z_3}=-\frac{du}{\alpha u+X_5}.$$

The integrals of the system are

$$u\exp\left(\int\frac{\alpha dz}{Z_3}\right)+\int\frac{X_5}{Z_3}\exp\left(\int\frac{\alpha\,dz}{Z_3}\right)dz=\text{const},\quad x=\text{const},$$
$$y\exp\left(-\int\frac{\alpha+Z_3'}{Z_3}\,dz\right)-\int\frac{Z_4}{Z_3}\exp\left(-\int\frac{\alpha+Z_3'}{Z_3}dz\right)dz=\text{const}\,.$$

We introduce new functions

$$Z=\int\frac{dz}{Z_3},\quad \bar{Z}=\int\frac{Z_4}{Z_3}\exp\left(-\int\frac{\alpha+Z_3'}{Z_3}\,dz\right)dz,\quad X=X_5.$$

Note that $Z'\neq 0$, We and again distinguish two subcases:

Subcase 01.0**:** $\alpha=0$;

Subcase 01.1**:** $\alpha\neq 0$.

Subcase 01.0. In this case, the system of integrals transforms to

$$u+XZ=\text{const},\quad x=\text{const},\quad yZ'-\bar{Z}=const,$$

and thus

$$\Psi(u+XZ,x,yZ'-\bar{Z})=0 \tag{15}$$

is the general solution. Consequently,

$$u=\Phi(x,\xi)-XZ,$$

where $\xi=yZ'-\bar{Z}$. Substituting the last expression to Equation (10), we obtain the equation

$$\boxed{\Phi_{\xi\xi}=X'\Phi_\xi-X\Phi_{x\xi}.} \tag{16}$$

When $X=0$, we obtain the solutions

$$\boxed{u=a_1(yZ'-\bar{Z})+a_0,\qquad a_i=a_i(x).}$$

If $X\neq 0$ Equation (16) can also be solved and the general solution is

$$\boxed{u=\Phi\left(yZ'-\bar{Z}+\int\frac{dx}{X}\right)+a(x),}$$

where Φ is an arbitrary function in one argument.

Subcase 01.1. We can set $\alpha=1$ and the general solution is

$$\Psi((u+X)e^Z,x,Z'e^{-Z}y-\bar{Z})=0.$$

Hence, after the change $Z\mapsto\ln Z$, we have

$$u=\frac{1}{Z}\Phi(x,\xi)-X, \tag{17}$$

where $\xi=Z'y/Z^2-\bar{Z}$. Substituting (17) to Equation (10), one obtains

$$\boxed{\Phi_{\xi\xi}=\Phi_x\Phi_\xi-\Phi\Phi_{x\xi}.}$$

2.2.3. *Case* 10. System (12) is now of the form

$$\frac{dx}{X_2}=\frac{dy}{\alpha y+Z_4}=\frac{dz}{0}=-\frac{du}{(\alpha-X_2')u+X_5}.$$

Then the integrals are

$$u\exp\left(\int\frac{\alpha-X_2'}{X_2}\,dx\right)+\int\frac{X_5}{X_2}\exp\left(\int\frac{\alpha-X_2'}{X_2}\,dx\right)dx=\text{const},$$
$$y\exp\left(-\int\frac{\alpha\,dx}{X_2}\right)-\int\frac{Z_4}{X_2}\exp\left(-\int\frac{\alpha\,dx}{X_2}\right)dx=\text{const},\ z=\text{const}\,.$$

Let us introduce the notation

$$\int\frac{dx}{X_2}=X,\quad \frac{X_5}{X_2}\exp\left(\int\frac{\alpha-X_2'}{X_2}\,dx\right)dx=\bar{X},\quad Z_4=Z$$

and consider the subcases

Subcase 10.0**:** $\alpha=0$;

Subcase 10.1**:** $\alpha \neq 0$.

Subcase 10.0. In this case the general solution is given by

$$\Psi(X'u + \bar{X}, y - XZ, z) = 0, \qquad X' \neq 0,$$

and thus

$$u = \frac{1}{X'}\Phi(\xi, z) - \bar{X},$$

where $\xi = y - XZ$. Substituting this expression to Equation (10), one obtains

$$\boxed{(1 + Z\Phi_z)\Phi_{\xi\xi} = Z\Phi_\xi\Phi_{\xi z} + Z'\Phi_\xi^2.}$$

The equation can be solved explicitly. Indeed, dividing by Φ_ξ^2 one obtains

$$\frac{\Phi_{\xi\xi}}{\Phi_\xi^2} - Z' = Z\frac{\Phi_\xi\Phi_{\xi z} - \Phi_z\Phi_{\xi\xi}}{\Phi_\xi^2},$$

or

$$-\left(\frac{1}{\Phi_\xi}\right)_\xi - Z' = Z\left(\frac{\Phi_z}{\Phi_\xi}\right)_\xi.$$

Hence,

$$-\frac{1}{\Phi_\xi} - Z'\xi = Z\frac{\Phi_z}{\Phi_\xi} + \varphi,$$

where $\varphi = \varphi(z)$ is an arbitrary function. Thus,

$$Z\Phi_z + (Z'\xi + \varphi)\Phi_\xi = -1$$

and

$$\boxed{\Phi = \Upsilon(\xi - \bar{\varphi}) - \int \frac{dz}{Z}}$$

is the general solution, where $\bar{\varphi} = Z\int \frac{\varphi\, dz}{Z^2}$.

Subcase 10.1. We may set $\alpha = 1$ and then obtain the general solution in the form

$$\Psi\left(X'ue^X + \bar{X}, e^{-X}(y + Z), z\right) = 0,$$

or, after the change $X \mapsto \ln X$,

$$u = \frac{\Phi(\xi, z) - \bar{X}}{X'}, \tag{18}$$

where $\xi = (y + Z)/X$. Substituting to (10), one has

$$\boxed{(1 + \xi\Phi_z)\Phi_{\xi\xi} - \xi\Phi_\xi\Phi_{\xi z} + \Phi_\xi\Phi_z = 0.}$$

2.2.4. *Case* 11. We have here

$$\frac{dx}{X_2} = \frac{dy}{(\alpha + Z_3')y + Z_4} = \frac{dz}{Z_3} = -\frac{du}{(\alpha - X_2')u + X_5},$$

where $X_2 \neq 0$, $Z_3 \neq 0$. The integrals are

$$u\exp\left(\int \frac{\alpha - X_2'}{X_2}\,dx\right) + \int \frac{X_5}{X_2}\exp\left(\int \frac{\alpha - X_2'}{X_2}\,dx\right)dx = \text{const}, \int \frac{dx}{X_2} - \int \frac{dx}{Z_3} = \text{const},$$

$$y\exp\left(-\int \frac{\alpha + Z_3'}{Z_3}\,dx\right) - \int \frac{Z_4}{Z_3}\exp\left(-\int \frac{\alpha + Z_3'}{Z_3}\,dx\right)dx = \text{const}\,.$$

As before, we introduce the notation

$$\int \frac{dx}{X_2} = X, \ \int \frac{X_5}{X_2}\exp\left(\int \frac{\alpha - X_2'}{X_2}\,dx\right)dx = \bar{X}, \quad \int \frac{dx}{Z_3} = Z, \ \int \frac{Z_4}{Z_3}\exp\left(-\int \frac{\alpha + Z_3'}{Z_3}\,dx\right)dx = \bar{Z}$$

and consider two subcases

Subcase 11.0**:** $\alpha = 0$;
Subcase 11.1**:** $\alpha \neq 0$.

Subcase 11.0. The general solution is given by

$$\Psi(X'u+\bar{X}, Z'y-\bar{Z}, X-Z)=0$$

in this case and thus

$$u=\frac{\Phi(\xi,\eta)-\bar{X}}{X'}, \qquad \xi=Z'y-\bar{Z}, \quad \eta=X-Z.$$

After substituting to Equation (10), one has

$$\boxed{\Phi_{\xi\xi}=\Phi_\xi\Phi_{\eta\eta}-\Phi_\eta\Phi_{\xi\eta}.}$$

The equation can be linearized by the Legendre transformation, [12].

Subcase 11.1. Setting $\alpha=1$, we obtain the general solution

$$\Psi(X'e^X u+\bar{X}, Z'e^{-Z}y-\bar{Z}, X-Z)=0$$

and, after the change $X\mapsto \ln X$, $Z\mapsto -\ln Z$, one has

$$u=\frac{\Phi(\xi,\eta)-\bar{X}}{X'}, \qquad \xi=Z'y-\bar{Z}, \quad \eta=\ln(XZ). \tag{19}$$

Substituting (19) to (10), we obtain the equation

$$\boxed{\Phi_\eta\Phi_{\xi\eta}-\Phi_\xi\Phi_{\eta\eta}=e^\eta\Phi_{\xi\xi}.}$$

3. The 3D rdDym equation

The equation is

$$u_{ty}=u_xu_{xy}-u_yu_{xx}. \tag{20}$$

3.1. **Symmetries.** Symmetries of Equation (20) are defined by

$$D_tD_y(\varphi)=u_xD_xD_y(\varphi)-u_yD_x^2(\varphi)+u_{xy}D_x(\varphi)-u_{xx}D_y(\varphi). \tag{21}$$

Solutions of (21) are

$$\begin{aligned}
\varphi_1&=xu_x-2u,\\
\varphi_2(T_2)&=T_2u_t+T_2'(xu_x-u)+\frac{1}{2}T_2''x^2,\\
\varphi_3(Y_3)&=Y_3u_y,\\
\varphi_4(T_4)&=T_4u_x+T_4'x,\\
\varphi_5(T_5)&=T_5,
\end{aligned}$$

where $Y_i=Y_i(y)$, $T_i=T_i(t)$ and 'primes' denote the derivatives. The commutator relations are given in Table 3.

	φ_1	$\varphi_2(\bar{T}_2)$	$\varphi_3(\bar{Y}_3)$	$\varphi_4(\bar{T}_4)$	$\varphi_5(\bar{T}_5)$
φ_1	0	0	0	$\varphi_4(T_4)$	$2\varphi_5(T_5)$
$\varphi_2(T_2)$	...	$\varphi_2(\bar{T}_2X_2'-T_2\bar{T}_2')$	0	$\varphi_4(\bar{T}_4T_2'-T_2\bar{T}_4')$	$\varphi_5(\bar{T}_5T_2'-T_2\bar{T}_5')$
$\varphi_3(Y_3)$	...	...	0	0	0
$\varphi_4(T_4)$	...	...	...	$\varphi_5(\bar{T}_4T_4'-T_4\bar{T}_4')$	0
$\varphi_5(T_5)$	...	...	...	...	0

Table 3. Lie algebra structure of $\operatorname{sym}\mathscr{E}_{(20)}$

3.2. **Reductions.** The general symmetry of Equation (20) is

$$\varphi=(\alpha x+T_2'x+T_4)u_x+Y_3u_y+T_2u_t-(2\alpha+T_2')u+T_5+T_4'x+\frac{1}{2}T_2''x^2,$$

where $\alpha\in\mathbb{R}$ is a constant. Consequently, φ-invariant solutions are defined by the system

$$\frac{dx}{(\alpha+T_2')x+T_4}=\frac{dy}{Y_3}=\frac{dt}{T_2}=\frac{du}{(2\alpha+T_2')u-T_5-T_4'x-\frac{1}{2}T_2''x^2}. \tag{22}$$

In what follows, we consider the following cases

Case 00**:** $Y_3=0$, $T_2=0$;
Case 01**:** $Y_3=0$, $T_2\neq 0$;
Case 10**:** $Y_3\neq 0$, $T_2=0$;
Case 11**:** $Y_3\neq 0$, $T_2\neq 0$.

3.2.1. *Case* 00. Equation (29) takes the form

$$\frac{dx}{\alpha x + T_4} = \frac{dy}{0} = \frac{dt}{0} = \frac{du}{2\alpha u - T_5 - T_4' x}.$$

As before, two subcase must be considered:

Subcase 00.0**:** $\alpha = 0$;
Subcase 00.1**:** $\alpha \neq 0$.

Subcase 00.0. Here we have

$$\frac{dx}{T_4} = \frac{dy}{0} = \frac{dt}{0} = -\frac{du}{T_5 + T_4' x}$$

and the general solution is given by

$$\Psi\left(u + \frac{1}{2}Tx^2 + \bar{T}x, y, t\right) = 0,$$

or

$$u = \Phi(y,t) - \frac{1}{2}Tx^2 - \bar{T}x, \tag{23}$$

where $T = T_4'/T_4$, $\bar{T} = T_5/T_4$. Substituting (23) in Equation (19), we obtain

$$\boxed{\Phi_{yt} = T\Phi_y.}$$

The general solution is

$$\Phi = \varphi(y)e^{\int T\,dt} + \psi(t),$$

which leads to the following family of solutions to Equation (19):

$$\boxed{u = \varphi(y)e^{\int T\,dt} + \psi(t) - \frac{1}{2}Tx^2 - \bar{T}x.}$$

Subcase 00.1. Setting $\alpha = 1$, we obtain

$$\frac{dx}{x + T_4} = \frac{dy}{0} = \frac{dt}{0} = \frac{du}{2u - T_5 - T_4' x}.$$

The general solution of this system is

$$u = (x+T)^2\Phi(y,t) + T'(x+T) + \bar{T}, \tag{24}$$

where $T = T_4$, $\bar{T} = (T_5 - T_4T_4')/2$. Substituting to (20), we obtain the equation

$$\boxed{\Phi_{yt} = 2\Phi\Phi_y.}$$

Integrating over y, we come to the Riccati equation

$$\Phi_t = \Phi^2 + \varphi(t).$$

Thus, to any choice of φ there corresponds a family of solutions to Equation (20).

Examples. Let us consider some particular cases.

(1) If $\varphi = 0$ then

$$\Phi = \frac{1}{\psi - t}.$$

Here and in all the examples below ψ is an arbitrary function of y.

(2) For $\varphi = a^2$, $a = \text{const}$, one has

$$\Phi = a\tan\big(a(t+\psi)\big).$$

(3) If $\varphi = -a^2$ then

$$\Phi = a\frac{1 + e^{2a(t+\psi)}}{1 - e^{2a(t+2\psi)}}$$

(4) For $\varphi = t^\varkappa$ one has

$$\Phi = \frac{-\psi J(\frac{1}{\varkappa+2}, \frac{2t^{\frac{1}{2}\varkappa+1}}{\varkappa+2}) + \psi J(\frac{3+\varkappa}{\varkappa+2}, \frac{2t^{\frac{1}{2}\varkappa+1}}{\varkappa+2})t^{\frac{1}{2}\varkappa+1} - Y(\frac{1}{\varkappa+2}, \frac{2t^{\frac{1}{2}\varkappa+1}}{\varkappa+2}) + Y(\frac{3+\varkappa}{\varkappa+2}, 2\frac{t^{\frac{1}{2}\varkappa+1}}{\varkappa+2})t^{\frac{1}{2}\varkappa+1}}{t\big(\psi J(\frac{1}{\varkappa+2}, \frac{2t^{\frac{1}{2}\varkappa+1}}{\varkappa+2}) + Y(\frac{1}{\varkappa+2}, \frac{2t^{\frac{1}{2}\varkappa+1)}}{\varkappa+2})\big)},$$

where

$$J(a,\theta) = \sum_{m=0}^{\infty} \frac{(-1)^m}{m!\,\Gamma(m+a+1)}\left(\frac{\theta}{2}\right)^{2m+a},$$

$$Y(a,\theta) = \frac{J(a,\theta)\cos(a\pi) - J(-a,\theta)}{\sin(a\pi)}$$

are Bessel functions of the first and second kinds, respectively.

(5) If $\varphi = e^t$, then

$$\Phi = \left(\frac{\psi Y(1, 2e^{\frac{t}{2}})}{\psi Y(0, 2e^{\frac{t}{2}}) + J(0, 2e^{\frac{t}{2}})} + \frac{J(1, 2e^{\frac{t}{2}})}{\psi Y(0, 2e^{\frac{t}{2}}) + J(0, 2e^{\frac{t}{2}})} \right) e^{\frac{t}{2}}$$

(6) For $\varphi = (1-t)/(1+t)$ the solution is

$$\Phi = \frac{2\psi \,\mathrm{Ei}(1, -2-2t)t + \psi e^{2+2t} + 2t}{2\psi(1+t)\,\mathrm{Ei}(1, -2-2t) + 2t + \psi e^{2+2t} + 2},$$

where

$$\mathrm{Ei}(a, t) = \int_1^\infty \frac{e^{-\theta t}\, d\theta}{\theta^a}$$

is the exponential integral function.

3.2.2. *Case* 01. We have

$$\frac{dx}{(\alpha + T_2')x + T_4} = \frac{dy}{0} = \frac{dt}{T_2} = \frac{du}{(2\alpha + T_2')u - T_5 - T_4' x - \frac{1}{2}T_2'' x^2},$$

where $T_2 \neq 0$. Its integrals are

$$T' x e^{-\alpha T} - \bar{T} = \text{const}, \; y = \text{const},$$

$$T' u e^{-2\alpha T} + \bar{\bar{T}} + \left(\alpha \bar{T} + \left(\frac{\bar{T}}{T'} \right)' \right) (T' x e^{-\alpha T} - \bar{T}) - \frac{1}{2} \frac{T''}{(T')^2} (T' x e^{-\alpha T} - \bar{T})^2 = \text{const},$$

where

$$T = \int \frac{dt}{T_2}, \quad \bar{T} = \int \left(T_4 \cdot (T')^2 \cdot e^{-\alpha T} \right) \, dt,$$

$$\bar{\bar{T}} = \int \left(T_5 \cdot (T')^2 \cdot e^{-2\alpha T} + T_4' \cdot T \cdot T' \cdot e^{-\alpha T} + \frac{1}{2} T_2'' \cdot \bar{T}^2 \right) \, dt.$$

Then the general solution is

$$\Psi \left(T' x e^{-\alpha T} - \bar{T}, y, T' u e^{-2\alpha} T + \bar{\bar{T}} + \left(\alpha \bar{T} + \left(\frac{\bar{T}}{T'} \right)' \right) (T' x e^{-\alpha T} - \bar{T}) \right.$$

$$\left. - \frac{1}{2} \frac{T''}{(T')^2} (T' x e^{-\alpha T} - \bar{T})^2 \right) = 0,$$

or

$$u = \left(\Phi(\xi, y) - \bar{\bar{T}} - \left(\alpha \bar{T} + \left(\frac{\bar{T}}{T'} \right)' \right) \xi + \frac{1}{2} \frac{T''}{(T')^2} \xi^2 \right) \frac{e^{2\alpha T}}{T'},$$

where

$$\xi = T' x e^{-\alpha T} - \bar{T}.$$

Substituting to Equation (20), one obtains

$$\boxed{(\alpha \xi + \Phi_\xi)\Phi_{\xi y} - \Phi_y(\Phi_{\xi\xi} + 2\alpha) = 0.} \tag{25}$$

3.2.3. *Case* 10. The defining equations are

$$\frac{dx}{\alpha x + T_4} = \frac{dy}{Y_3} = \frac{dt}{0} = \frac{du}{2\alpha u - T_5 - T_4' x}, \tag{26}$$

where $Y_3 \neq 0$. Below we consider the following subcases:

Subcase 10.00**:** $\alpha = 0$, $T_4 = 0$;
Subcase 10.01**:** $\alpha = 0$, $T_4 \neq 0$;
Subcase 10.1**:** $\alpha \neq 0$

and introduce $Y = Y(y)$ such that $1/Y_3 = Y'$.

Subcase 10.00. In this case, $T_5 \neq 0$ and System (26) takes the form

$$\frac{dx}{0} = Y'\,dy = \frac{dt}{0} = -\frac{du}{T_5}.$$

Denote $T_5 = T$. Then the integrals are

$$x = \text{const}, \quad t = \text{const}, \quad u + YT = \text{const}\,.$$

Then the general solution is given by

$$\Psi(u + YT, x, t) = 0,$$

or

$$u = \Phi(x, t) - YT.$$

Substituting to Equation (20), one obtains

$$-Y'T' = Y'T\Phi_{xx},$$

or, since $Y' = 1/Y_3 \neq 0$,

$$\Phi_{xx} = -\frac{T'}{T}.$$

This delivers us the following family of solutions:

$$\boxed{u = -\frac{T'}{2T}x^2 + \varphi(t)x + \psi(t) - YT.}$$

Subcase 10.01. The defining equations are now

$$\frac{dx}{T_4} = Y'\,dy = \frac{dt}{0} = -\frac{du}{T_5 + T_4'x}.$$

Let us introduce the notation $T_4 = T$, $T_5/T_4 = \bar{T}$. Then the integrals are

$$x - YT = \text{const}, \quad t = \text{const}, \quad u + \frac{T'}{2T}x^2 + \bar{T}x = \text{const}$$

and the general solution is

$$\Psi(u + \frac{T'}{2T}x^2 + \bar{T}x, x - YT, t) = 0,$$

or

$$u = \Phi(\xi, t) - \frac{T'}{2T}x^2 - \bar{T}x,$$

where $\xi = x - YT$. Substituting to (20), we obtain the linear equation

$$\boxed{\left(\frac{T'}{T}\xi + \bar{T}\right)\Phi_{\xi\xi} + \Phi_{\xi t} = 0.}$$

The general solution of this equation is

$$\Phi = \varphi(\eta)T + \psi(t), \qquad \eta = \frac{\xi}{T} - \int \frac{\bar{T}}{T}\,dt,$$

which gives the family of solutions to (20):

$$\boxed{u = \varphi(\eta)T + \psi(t) - \frac{T'}{2T}x^2 - \bar{T}x, \qquad \eta = \frac{\xi}{T} - \int \frac{\bar{T}}{T}\,dt.}$$

Subcase 10.1. We can assume $\alpha = 1$ and the defining equations become

$$\frac{dx}{x + T_4} = Y'\,dy = \frac{dt}{0} = \frac{du}{2u - T_5 - T_4'x}.$$

The integrals of this system are

$$(x + T)e^{-Y} = \text{const}, \quad t = \text{const}, \quad \frac{u - \bar{T}}{(x + T)^2} - \frac{T'}{x + T} = \text{const},$$

where $T = T_4$, $\bar{T} = (T_5 - T'T)/2$, and thus the general solution is

$$\Psi\left(\frac{u - \bar{T}}{(x + T)^2} - \frac{T'}{x + T}(x + T), e^{-Y}, t\right) = 0,$$

or

$$u = (x + T)^2\Phi(\xi, t) + T'(x + T) + \bar{T}, \qquad \xi = (x + T)e^{-Y}.$$

Substituting to (20), one obtains the equation

$$\boxed{\Phi_{\xi t} = 4\Phi\Phi_\xi - \xi\Phi_\xi^2 + 2\xi\Phi\Phi_{\xi\xi}.}$$

3.2.4. *Case* 11. Let us set $Y' = 1/Y_3 \neq 0$ and $T' = 1/T_2 \neq$. Then System (22) becomes

$$\frac{dx}{(\alpha + T_2')x + T_4} = Y'\,dy = T'\,dt = \frac{du}{(2\alpha + T_2')u - T_5 - T_4'x - \frac{1}{2}T_2''x^2}.$$

The integrals are

$$T'xe^{-\alpha T} - \bar{T} = \text{const}, \quad Y - T = \text{const},$$

$$T'ue^{-2\alpha} + \bar{\bar{T}} + \left(\alpha\bar{T} + \left(\frac{\bar{T}}{T'}\right)'\right)(T'xe^{-\alpha T} - \bar{T}) - \frac{1}{2}\frac{T''}{(T')^2}(T'xe^{-\alpha T} - \bar{T})^2 = \text{const},$$

where, as before,

$$T = \int \frac{dt}{T_2}, \quad \bar{T} = \int \left(T_4 \cdot (T')^2 \cdot e^{-\alpha T}\right)\,dt,$$

$$\bar{\bar{T}} = \int \left(T_5 \cdot (T')^2 \cdot e^{-2\alpha T} + T_4' \cdot T \cdot T' \cdot e^{-\alpha T} + \frac{1}{2}T_2'' \cdot \bar{T}^2\right)\,dt.$$

Thus, the general solution is given by

$$\Psi\left(T'xe^{-\alpha T} - \bar{T}, Y - T, T'ue^{-2\alpha}T + \bar{\bar{T}} + \left(\alpha\bar{T} + \left(\frac{\bar{T}}{T'}\right)'\right)(T'xe^{-\alpha T} - \bar{T})\right.$$

$$\left. -\frac{1}{2}\frac{T''}{(T')^2}(T'xe^{-\alpha T} - \bar{T})^2\right) = 0,$$

or

$$u = \left(\Phi(\xi,\eta) - \bar{\bar{T}} - \left(\alpha\bar{T} + \left(\frac{\bar{T}}{T'}\right)'\right)\xi + \frac{1}{2}\frac{T''}{(T')^2}\xi^2\right)\frac{e^{2\alpha T}}{T'},$$

where

$$\xi = T'xe^{-\alpha T} - \bar{T}, \qquad \eta = Y - T.$$

Substituting the last expression to Equation (20), one obtains

$$\boxed{\Phi_{\eta\eta} + (\alpha\xi + \Phi_\eta)\Phi_{\xi\eta} = \Phi_\eta(2\alpha + \Phi_{\xi\xi}).}$$

4. The basic Veronese web equation

The equation is

$$u_{ty} = u_t u_{xy} - u_y u_{tx}. \tag{27}$$

4.1. **Symmetries.** Symmetries of (27) are defined by

$$D_t D_y(\varphi) = u_t D_x D_y(\varphi) - u_y D_t D_x(\varphi) + u_{xy} D_t(\varphi) - u_{tx} D_y(\varphi), \tag{28}$$

whose solutions are

$$\begin{aligned}
\varphi_1(T_1) &= T_1 u_t,\\
\varphi_2(X_2) &= X_2 u_x - X_2' u,\\
\varphi_3(Y_3) &= Y_3 u_y,\\
\varphi_4(X_4) &= X_4,
\end{aligned}$$

where $X_i = X_i(x)$, $Y_i = Y_i(y)$, and $T_i = T_i(t)$. The commutator relations in $\operatorname{sym}\mathscr{E}_{(27)}$ are given in Table 4.

	$\varphi_1(\bar{T}_1)$	$\varphi_2(\bar{X}_2)$	$\varphi_3(\bar{Y}_3)$	$\varphi_4(\bar{X}_4)$
$\varphi_1(T_1)$	$\varphi_1(\bar{T}_1T_1' - T_1\bar{T}_1')$	0	0	0
$\varphi_2(X_2)$	...	$\varphi_2(\bar{X}_2X_2' - X_2\bar{X}_2')$	0	$\varphi_4(\bar{X}_4X_2' - X_2\bar{X}_4')$
$\varphi_3(Y_3)$	...	...	$\varphi_3(\bar{Y}_3Y_3' - Y_3\bar{Y}_3')$	0
$\varphi_4(X_4)$	...	...	...	0

Table 4. Lie algebra structure of $\operatorname{sym}\mathscr{E}_{(27)}$

4.2. **Reductions.** The general symmetry of Equation (27) is

$$\varphi = X_2 u_x + Y_3 u_y + T_1 u_t - X_2' u + X_4$$

and the corresponding invariant solutions must satisfy the system

$$\frac{dx}{X_2} = \frac{dy}{Y_3} = \frac{dt}{T_1} = \frac{du}{X_2' u - X_4}. \tag{29}$$

We consider below the following cases:

Case 100**:** $X_2 \neq 0$, $Y_3 = 0$, $Z_4 = 0$;
Case 010**:** $X_2 = 0$, $Y_3 \neq 0$, $Z_4 = 0$;
Case 001**:** $X_2 = 0$, $Y_3 = 0$, $Z_4 \neq 0$;
Case 011**:** $X_2 = 0$, $Y_3 \neq 0$, $Z_4 \neq 0$;
Case 101**:** $X_2 \neq 0$, $Y_3 = 0$, $Z_4 \neq 0$;
Case 110**:** $X_2 \neq 0$, $Y_3 \neq 0$, $Z_4 = 0$;
Case 111**:** $X_2 \neq 0$, $Y_3 \neq 0$, $Z_4 \neq 0$;

and use the notation $1/X_2 = X'$, $1/Y_3 = Y'$, $1/Z_4 = Z'$ when it is well defined.

4.2.1. *Case* 100. The defining equation is

$$\frac{dx}{X_2} = \frac{dy}{0} = \frac{dt}{0} = \frac{du}{X_2' u - X_4}.$$

The integrals are

$$X u - \bar{X} = \text{const}, \qquad y = \text{const}, \qquad t = \text{const},$$

and thus

$$\Psi(X u - \bar{X}, y, t) = 0$$

is the general solution, where $\bar{X} = \int X_4 X'\, dx$. Consequently,

$$u = \frac{\Phi(y,t) + \bar{X}}{X}.$$

Substituting to (27), one obtains

$$\boxed{\Phi_{yt} = 0.}$$

Hence, $\Phi = \varphi(y) + \psi(t)$ and

$$\boxed{u = \frac{\varphi(y) + \psi(t) + \bar{X}}{X}}$$

is a family of solutions to (27).

4.2.2. *Case* 010. The defining equation is

$$\frac{dx}{0} = \frac{dy}{Y_3} = \frac{dt}{0} = -\frac{du}{X_4}.$$

The integrals are

$$u + \bar{X} Y = \text{const}, \quad x = \text{const}, \quad t = \text{const},$$

where $\bar{X} = X_4$. Then

$$\Psi(u + \bar{X} Y, x, t) = 0$$

is the general solution and

$$u = \Phi(x,t) - \bar{X} Y.$$

Substituting to (27), one obtains $Y'(\bar{X}\Phi_{xt} - \bar{X}'\Phi_t) = 0$ and since $Y' \neq 0$,

$$\boxed{\bar{X}\Phi_{xt} - \bar{X}'\Phi_t = 0.}$$

Thus, if $\bar{X} = 0$ we obtain the obvious family of solutions

$$\boxed{u = \Phi(x,t).}$$

If $\bar{X} \neq 0$ the corresponding family of solutions is

$$\boxed{u = \bar{X}\varphi(t) + \psi(x) - \bar{X} Y.}$$

4.2.3. *Case* 001. The defining equation is

$$\frac{dx}{0} = \frac{dy}{0} = \frac{dt}{T_1} = -\frac{du}{X_4}.$$

Then, again denoting $\bar{X} = X_4$, we get the integrals

$$u + \bar{X}T = \text{const}, \quad x = \text{const}, \quad y = \text{const}$$

and the general solution in the form

$$\Psi(u + \bar{X}T, x, y) = 0,$$

or

$$u = \Phi(x, y) - \bar{X}T.$$

Substituting to (27), one obtains

$$\boxed{\bar{X}\Phi_{xy} - \bar{X}'\Phi_y = 0,}$$

since $T' \neq 0$. Then in the case $\bar{X} = 0$ we get the family of solutions

$$\boxed{u = \Phi(x, y)}$$

and when $\bar{X} \neq 0$ the family

$$\boxed{u = \bar{X}\varphi(y) + \psi(x) - \bar{X}T.}$$

4.2.4. *Case* 011. The defining equation is

$$\frac{dx}{0} = \frac{dy}{Y_3} = \frac{dt}{T_1} = -\frac{du}{X_4}.$$

Its integrals are

$$x = \text{const}, \quad Y - T = \text{const}, \quad u + \bar{X}Y = \text{const}$$

and the general solution is

$$\Psi(u + \bar{X}Y, x, Y - T) = 0,$$

or

$$u = \Phi(x, \xi) - \bar{X}Y, \qquad \xi = Y - T.$$

Substituting to Equation (27), one obtains

$$\boxed{\Phi_{\xi\xi} = \bar{X}\Phi_{x\xi} - \bar{X}'\Phi_\xi.}$$

If $\bar{X} = 0$ then

$$\boxed{u = \varphi(x) + \psi(Y - T) - \bar{X}Y.}$$

In the case $\bar{X} \neq 0$ the corresponding family is

$$\boxed{u = \bar{X}\varphi\left(Y - T + \int \frac{dx}{\bar{X}}\right) + \psi(x) - \bar{X}Y.}$$

4.2.5. *Case* 101. The defining equation is

$$\frac{dx}{X_2} = \frac{dy}{0} = \frac{dt}{T_1} = \frac{du}{X_2'u - X_4}.$$

The integrals of this system are

$$X'u + \bar{X} = \text{const}, \quad X - T = \text{const}, \quad y = \text{const},$$

where $\bar{X} = \int (X')^2 X_4\, dx$ and the general solution is given by

$$\Psi(X'u - \bar{X}, X - T, y) = 0,$$

or

$$u = \frac{\Phi(y, \xi) + \bar{X}}{X'}, \qquad \xi = X - T.$$

After substitution to Equation (27) one obtains

$$\boxed{(1 + \Phi_\xi)\Phi_{y\xi} = \Phi_y\Phi_{\xi\xi}.}$$

The general solution to this equation is

$$\Psi = \Upsilon\left(\xi + \psi(y)\right) - \xi$$

and thus we get the family

$$\boxed{u=\frac{\Upsilon\big(X-T+\psi(y)\big)-X+T+\bar{X}}{X'}}$$

to Equation (27).

4.2.6. *Case* 110. The defining equation is

$$\frac{dx}{X_2}=\frac{dy}{Y_3}=\frac{dt}{0}=\frac{du}{X_2'u-X_4}.$$

The integrals of this system are

$$X'u-\bar{X}=\text{const},\quad X-Y=\text{const},\quad y=\text{const},$$

where $\bar{X}=\int (X')^2X_4\,dx$ and the general solution is given by

$$\Psi(X'u+\bar{X},X-Y,y)=0,$$

or

$$u=\frac{\Phi(y,\xi)+\bar{X}}{X'},\qquad \xi=X-Y.$$

Substitution to Equation (27) leads to

$$\boxed{(1+\Phi_\xi)\Phi_{t\xi}=\Phi_t\Phi_{\xi\xi}.}$$

Similar to the previous case, we solve this equation and obtain the following family of solutions to Equation (27):

$$\boxed{u=\frac{\Upsilon\big(X-Y+\psi(t)\big)-X+Y+\bar{X}}{X'}.}$$

4.2.7. *Case* 111. The defining equation is

$$\frac{dx}{X_2}=\frac{dy}{Y_3}=\frac{dt}{T_1}=\frac{du}{X_2'u-X_4}.$$

The integrals are

$$X'u-\bar{X}=\text{const},\quad X-Y=\text{const},\quad X-T=\text{const},$$

where, as before $\bar{X}=\int (X')^2X_4\,dx$. This delivers the general solution

$$\Psi(X'u-\bar{X},X-Y,X-T)=0,$$

i.e.,

$$u=\frac{\Phi(\xi,\eta)+\bar{X}}{X'},\qquad \xi=X-Y,\quad \eta=X-T.$$

Substituting to (27), we obtain the equation

$$\boxed{\Phi_\eta\Phi_{\xi\xi}+(\Phi_\eta-\Phi_\xi-1)\,\Phi_{\xi\eta}-\Phi_\xi\Phi_{\eta\eta}=0.}\tag{30}$$

The equation linearizes by the Legendre transformation, [12].

5. Pavlov's equation

The equation is

$$u_{yy}=u_{tx}+u_yu_{xx}-u_xu_{xy}.\tag{31}$$

5.1. **Symmetries.** The defining equation for symmetries of (31) is

$$D_y^2(\varphi)=D_tD_x(\varphi)+u_yD_x^2(\varphi)-u_xD_xD_y(\varphi)+u_{xx}D_y(\varphi)-u_{xy}D_x(\varphi).\tag{32}$$

Its solutions are

$$\begin{aligned}
\varphi_1&=2x-yu_x,\\
\varphi_2&=3u-2xu_x-yu_y,\\
\varphi_3(T_3)&=T_3u_t+T_3'(xu_x+yu_y-u)+\frac{1}{2}T_3''(y^2u_x-2xy)-\frac{1}{6}T_3'''y^3,\\
\varphi_4(T_4)&=T_4u_x-T_4'y,\\
\varphi_5(T_5)&=T_5u_y+T_5'(yu_x-x)-\frac{1}{2}T_5''y^2,\\
\varphi_6(T_6)&=T_6,
\end{aligned}$$

where T_i are functions of t. The Lie algebra structure in $\operatorname{sym}\mathscr{E}_{(31)}$ is given in Table 5.

	φ_1	φ_2	$\varphi_3(\bar{T}_3)$	$\varphi_4(\bar{T}_4)$	$\varphi_5(\bar{T}_5)$	$\varphi_6(\bar{T}_6)$
φ_1	0	φ_1	0	$2\varphi_6(T_4)$	$-2\varphi_4(T_5)$	0
φ_2	...	0	0	$-2\varphi_4(\bar{T}_4)$	$-\varphi_5(\bar{T}_5)$	$-3\varphi_6(\bar{T}_6)$
$\varphi_3(T_3)$	...	...	$\varphi_3(\bar{T}_3T_3' - T_3\bar{T}_3')$	$\varphi_4(\bar{T}_4T_3' - T_3\bar{T}_4')$	$\varphi_5(\bar{T}_5T_3' - T_3\bar{T}_5')$	$\varphi_6(\bar{T}_6T_3 - T_3\bar{T}_6')$
$\varphi_4(T_4)$	...	...	...	0	$\varphi_6(\bar{T}_5T_4' - T_4\bar{T}_5')$	0
$\varphi_5(T_5)$	...	...	...	...	$\varphi_4(\bar{T}_5T_5' - T_5\bar{T}_5')$	0
$\varphi_6(T_6)$	...	...	...	...	...	0

TABLE 5. Lie algebra structure of $\operatorname{sym}\mathscr{E}_{(31)}$

5.2. **Reductions.** The general symmetry of Equation (31) is

$$\varphi = \left(-\alpha y - 2\beta x + T_3'x + \frac{1}{2}T_3''y^2 + T_4 + T_5'y\right)u_x + (-\beta y + T_3'y + T_5)u_y + T_3u_t + (3\beta - T_3')u + 2\alpha x - T_3''xy - \frac{1}{6}T_3'''y^3 - T_5'x - \frac{1}{2}T_5''y^2 + T_6 - T_4'y,$$

where α, $\beta \in \mathbb{R}$ are constants. Then the φ-invariant solutions are determined by the system

$$\frac{dx}{(T_3' - 2\beta)x + (T_5' - \alpha)y + \frac{1}{2}T_3''y^2 + T_4} = \frac{dy}{(T_3' - \beta)y + T_5} = \frac{dt}{T_3} = \frac{du}{(T_3' - 3\beta)u + (T_5' - 2\alpha)x + T_4'y + T_3''xy + \frac{1}{2}T_5''y^2 + \frac{1}{6}T_3'''y^3 - T_6}. \quad (33)$$

We consider the following tree of options:



5.2.1. *Case* 0. The defining equations are

$$\frac{dx}{-2\beta x + (T_5' - \alpha)y + T_4} = \frac{dy}{-\beta y + T_5} = \frac{dt}{0} = \frac{du}{-3\beta u + (T_5' - 2\alpha)x + T_4'y + \frac{1}{2}T_5''y^2 - T_6}.$$

Due to the above picture, consider the following subcases:

Subcase 00001**:** $\beta = 0$, $T_5 = 0$, $\alpha = 0$, $T_4 \neq 0$;
Subcase 0001**:** $\beta = 0$, $T_5 = 0$, $\alpha \neq 0$;
Subcase 001**:** $\beta = 0$, $T_5 \neq 0$;
Subcase 01**:** $\beta \neq 0$.

Subcase 00001. The defining equations are

$$\frac{dx}{T_4} = \frac{dy}{0} = \frac{dt}{0} = \frac{du}{T_4'y - T_6}.$$

Then the integrals are

$$u - (Ty - \bar{T})x = \text{const}, \qquad y = \text{const}, \qquad t = \text{const},$$

where $T = T_4'/T_4$, $\bar{T} = T_6/T_4$. Thus,

$$\Psi(u - (Ty - \bar{T})x, y, t) = 0$$

is the general solution and

$$u = \Phi(y, t) + (Ty - \bar{T})x.$$

Substituting to Equation (31), one obtains

$$\Phi_{yy} = (T' - T^2)y + T\bar{T} - \bar{T}',$$

which gives the family

$$\boxed{u = \frac{1}{6}(T' - T^2)y^3 + \frac{1}{2}(T\bar{T} - \bar{T}')y^2 + \varphi(t)y + \psi(t) + (Ty - \bar{T})x.}$$

of exact solutions to (31).

Subcase 0001. We may assume $\alpha=-1$ and the defining equations become

$$\frac{dx}{y+T_4}=\frac{dy}{0}=\frac{dt}{0}=\frac{du}{2x+T_4'y-T_6}.$$

Then the integrals are

$$(y+T_4)u-x^2-(T_4'y-T_6)x=\text{const},\quad y=\text{const},\quad t=\text{const}\,.$$

Consequently, the general solution is given by

$$\Psi\left((y+T_4)u-x^2-(T_4'y-T_6)x,y,t\right)=0$$

and thus

$$u=\frac{\Phi(y,t)+x^2+(T_4'y-T_6)x}{y+T_4},\tag{34}$$

or

$$u=\Phi(y,t)+T'x+\frac{x^2-\bar{T}x}{y+T},$$

where $T=T_4$, $\bar{T}=T_4T_4'+T_6$. After substituting to (31), we obtain the equation

$$\boxed{\Phi_{yy}=\frac{2\Phi_y}{y+T}+T''-\frac{\bar{T}'}{y+T}+\frac{\bar{T}^2}{(y+T)^3}.}$$

Solving this equation, we obtain the following family of solutions to Equation (31):

$$\boxed{u=\varphi(t)(y+T)^3-\frac{1}{2}T''(y+T)^2+\frac{1}{2}\bar{T}'(y+T)+\frac{2x^2-2\bar{T}x+\bar{T}^2}{y+T}+T'x+\psi(t).}$$

Subcase 001. The defining equations are

$$\frac{dx}{(T_5'-\alpha)y+T_4}=\frac{dy}{T_5}=\frac{dt}{0}=\frac{du}{(T_5'-2\alpha)x+T_4'y+\frac{1}{2}T_5''y^2-T_6}.$$

Let us introduce the notation $T=1/T_5$, $\bar{T}=T_4/T_5$, $\bar{\bar{T}}=T_6/T_5$. Then the integrals acquire the form

$$t=\text{const},\quad x+\frac{1}{2}\left(\frac{T'}{T}+\alpha T\right)y^2-\bar{T}y=\text{const},$$
$$u+\left(\frac{1}{6}\frac{T''}{T}+\alpha T'+\frac{2}{3}\alpha^2T^2\right)y^3-\frac{1}{2}\left(\bar{T}'+2\alpha T\bar{T}\right)y^2+\left(\left(\frac{T'}{T}+2\alpha T\right)x+\bar{\bar{T}}\right)y=\text{const}\,.$$

Hence, the general solution is

$$\Psi\left(u+\left(\frac{1}{6}\frac{T''}{T}+\alpha T'+\frac{2}{3}\alpha^2T^2\right)y^3-\frac{1}{2}\left(\bar{T}'+2\alpha T\bar{T}\right)y^2+\left(\left(\frac{T'}{T}+2\alpha T\right)x+\bar{\bar{T}}\right)y,\right.$$
$$\left.x+\frac{1}{2}\left(\frac{T'}{T}+\alpha T\right)y^2-\bar{T}y,t\right)=0,$$

or

$$u=\Phi(\xi,t)-\left(\frac{1}{6}\frac{T''}{T}+\alpha T'+\frac{2}{3}\alpha^2T^2\right)y^3+\frac{1}{2}\left(\bar{T}'+2\alpha T\bar{T}\right)y^2-\left(\left(\frac{T'}{T}+2\alpha T\right)x+\bar{\bar{T}}\right)y,$$

where

$$\xi=x+\frac{1}{2}\left(\frac{T'}{T}+\alpha T\right)y^2-\bar{T}y.\tag{35}$$

Substituting to Equation (31), one obtains

$$\boxed{\left(\left(\frac{T'}{T}+2\alpha T\right)\xi+\bar{T}^2+\bar{\bar{T}}\right)\Phi_{\xi\xi}-\Phi_{\xi t}-\alpha T\Phi_\xi+\bar{T}'+2\alpha T\bar{T}=0.}$$

Of course, the equation can be solved explicitly, but the final result is too cumbersome: it is easily shown that

$$\Phi_\xi=\left(\bar{Z}\left(\xi e^{\int a\,dt}-\int be^{\int a\,dt}\,dt\right)+\int ce^{\alpha\int T\,dt}\,dt\right)e^{-\alpha\int T\,dt},$$

where $\bar{Z}$ is an arbitrary function in one variable and

$$a=\frac{T'}{T}+2\alpha T,\quad b=\bar{T}^2+\bar{\bar{T}},\quad c=\bar{T}'+2\alpha T\bar{T}.$$

Thus,

$$\Phi=Z\left(\xi e^{\int a\,dt}-\int be^{\int a\,dt}\,dt\right)e^{-2\alpha\int T\,dt}+\left(\int ce^{\alpha\int T\,dt}\,dt\right)\xi e^{-\alpha\int T\,dt}+\varphi(t),$$

and the corresponding family of solutions is

$$\boxed{\begin{aligned} u = Z\left(\xi e^{\int a\,dt} - \int b e^{\int a\,dt}\,dt\right) e^{-2\alpha\int T\,dt} + \left(\int c e^{\alpha\int T\,dt}\,dt\right)\xi e^{-\alpha\int T\,dt} + \varphi(t) \\ - \left(\frac{1}{6}\frac{T''}{T} + \alpha T' + \frac{2}{3}\alpha^2 T^2\right) y^3 + \frac{1}{2}\left(\bar{T}' + 2\alpha T\bar{T}\right) y^2 - \left(\left(\frac{T'}{T} + 2\alpha T\right) x + \bar{\bar{T}}\right) y \end{aligned}}$$

with ξ given by (35).

Subcase 01. Since $\beta \neq 0$, we can set $\beta = -1$ and the defining equations become

$$\frac{dx}{2x + (T_5' - \alpha)y + T_4} = \frac{dy}{y + T_5} = \frac{dt}{0} = \frac{du}{3u + (T_5' - 2\alpha)x + T_4' y + \frac{1}{2}T_5'' y^2 - T_6}. \tag{36}$$

Let us introduce the notation $T = T_5$, $\bar{T} = T_4 - T_5(T_5' - \alpha)$. Then the integrals of (36) are

$$t = \text{const}, \qquad \frac{x + \frac{1}{2}\bar{T}}{(y+T)^2} + \frac{T' - \alpha}{y + T} = \text{const}$$

$$\frac{u + (x + \frac{1}{2}\bar{T})(T' - 2\alpha) + \frac{1}{3}\bar{\bar{T}}}{(y+T)^3} + \frac{(T'-\alpha)^2 + \frac{1}{2}\bar{T}'}{(y+T)^2} + \frac{1}{2}\frac{T''}{y+T} = \text{const},$$

where

$$\bar{\bar{T}} = -\frac{1}{2}(T' - 2\alpha)\bar{T} - \left(\bar{T}' + T'(T' - \alpha) + TT''\right)T + \frac{1}{2}T^2 T'' - T_6.$$

Consequently, the general solution is

$$\Psi\left(\frac{u + (x + \frac{1}{2}\bar{T})(T' - 2\alpha) + \frac{1}{3}\bar{\bar{T}}}{(y+T)^3} + \frac{(T'-\alpha)^2 + \frac{1}{2}\bar{T}'}{(y+T)^2} + \frac{1}{2}\frac{T''}{y+T},\ \frac{x + \frac{1}{2}\bar{T}}{(y+T)^2} + \frac{T' - \alpha}{y+T},\ t\right) = 0,$$

or

$$u = (y+T)^3\Phi(\xi, t) - \frac{1}{2}T''(y+T)^2 - \left((T'-\alpha)^2 + \frac{1}{2}\bar{T}'\right)(y+T) - (T' - 2\alpha)\left(x + \frac{1}{2}\bar{T}\right) - \frac{1}{3}\bar{\bar{T}},$$

where

$$\xi = \frac{x + \frac{1}{2}\bar{T}}{(y+T)^2} + \frac{T' - \alpha}{y+T}.$$

Substituting to Equation (31), one obtains

$$\boxed{(4\xi^2 - 3\Phi)\Phi_{\xi\xi} - \Phi_{\xi t} - 6\xi\Phi_\xi + \Phi_\xi^2 + 6\Phi = 0.}$$

5.2.2. *Case* 1. The defining equation is now

$$\frac{dx}{(T_3' - 2\beta)x + (T_5' - \alpha)y + \frac{1}{2}T_3'' y^2 + T_4} = \frac{dy}{(T_3' - \beta)y + T_5} = \frac{dt}{T_3}$$

$$= \frac{du}{(T_3' - 3\beta)u + (T_5' - 2\alpha)x + T_4' y + T_3'' xy + \frac{1}{2}T_5'' y^2 + \frac{1}{6}T_3''' y^3 - T_6}$$

and since $T_3 \neq 0$ we may set $T' = 1/T_3$. The integrals are

$$T' y e^{\beta T} - \bar{T} = \text{const},$$

$$T' x e^{2\beta T} + \frac{1}{2}T_3'\xi^2 - \left(\left(\frac{\bar{T}}{T'}\right)' - \beta\bar{T}' - \alpha k(\beta)\right)\xi - \bar{\bar{T}} = \text{const},$$

$$T' u e^{3\beta T} - T_{00} - T_{10}\xi - T_{01}\eta - T_{20}\xi^2 - T_{11}\xi\eta - T_{30}\xi^3 = \text{const},$$

where

$$\xi = T' y e^{\beta T} - \bar{T},$$

$$\eta = T' x e^{2\beta T} + \frac{1}{2}T_3'\xi^2 - \left(\left(\frac{\bar{T}}{T'}\right)' - \beta\bar{T}' - \alpha k(\beta)\right)\xi - \bar{\bar{T}},$$

$$\bar{T} = \int T_5 (T')^2 e^{\beta T}\,dt,$$

$$\bar{\bar{T}} = \int \left(T_4 (T')^2 e^{2\beta T} + (T_5' - \alpha)\bar{T}T' e^{\beta T} + \frac{1}{2}T_3''(\bar{T})^2\right) dt,$$

$$
\begin{aligned}
T_{00} &= \int \left(T_3''\bar{\bar{T}}\bar{T} + \frac{T_3'''\bar{T}^3}{6T'} + \left(T_4'\bar{T}T' + (T_5' - 2\alpha)T' + \frac{1}{2T_5''(\bar{T})^2} \right) e^{\beta T} - T_6(T')^2 e^{3\beta T} \right) dt,\\
T_{10} &= \int \left(\frac{T_3'''(\bar{T})^2}{2T'} + T_3'' \left(\bar{\bar{T}} + \bar{T} \left(\left(\frac{\bar{T}}{T'} \right)' - \alpha k(\beta) - \beta\bar{T} \right) \right) \right.\\
&\quad \left. + \left(T_5''\bar{T} + (T_5' - 2\alpha) \left(\left(\frac{\bar{T}}{T'} \right)' - \alpha k(\beta) - \beta\bar{T} \right) \right) e^{\beta T} + T_4'T'e^{2\beta T} \right) dt,\\
T_{01} &= \int \left(T_3'''\bar{T} + (T_5' - 2\alpha)T'e^{\beta T} \right) dt,\\
T_{20} &= \int \left(T_3'' \left(\frac{\bar{T}}{2T'} + \left(\frac{\bar{T}}{T'} \right)' - \alpha k(\beta) - \beta\bar{T} - \frac{1}{2}T_3'\bar{T} \right) + \frac{1}{2}\left(T_5'' - (T_5' - 2\alpha)T' \right) e^{\beta T} \right) dt,\\
T_{11} &= \int T_3''\, dt = T_3',\\
T_{30} &= \int \left(\frac{T_3'''}{6T'} - \frac{1}{2}T_3''T_3' \right) dt = \frac{1}{6}T_3''T_3 - \frac{1}{3}(T_3')^2,
\end{aligned}
$$

and

$$
k(\beta) = \int T'e^{\beta T}\, dt = \begin{cases} \dfrac{e^{\beta T}}{\beta}, & \beta \neq 0,\\ T, & \beta = 0. \end{cases}
$$

Thus, the general solution is

$$
u = \left(\frac{\Phi(\xi,\eta) + T_{00} + T_{10}\xi + T_{01}\eta + T_{20}\xi^2 + T_{11}\xi\eta + T_{30}\xi^3}{T'} \right) e^{3\beta T}. \tag{37}
$$

Substituting (37) to Equation (31), one obtains

$$
\Phi_{\xi\xi} = (\beta\xi - \Phi_\eta)\,\Phi_{\xi\eta} + (2\beta\eta + \alpha\varkappa + \Phi_\xi)\,\Phi_{\eta\eta} - \beta\Phi_\varkappa - 2\alpha\varkappa,
$$

where

$$
\varkappa = e^{\beta T} - \beta k(\beta),
$$

i.e.,

$$
\varkappa = \begin{cases} 0, & \beta \neq 0,\\ \xi, & \beta = 0. \end{cases}
$$

Thus, the reductions are

$$
\boxed{\Phi_{\xi\xi} = (\beta\xi - \Phi_\eta)\Phi_{\xi\eta} + (2\beta\eta + \Phi_\xi)\Phi_{\eta\eta} - \beta\Phi_\eta = 0}
$$

for $\beta \neq 0$ and

$$
\boxed{\Phi_{\xi\xi} = (\alpha\xi + \Phi_\xi)\Phi_{\eta\eta} - \Phi_\eta\Phi_{\xi\eta} - 2\alpha} \tag{38}
$$

for $\beta = 0$. Note that in the case $\alpha \neq 0$ Equation (38) transforms to the Gibbons-Tsarev equation (see [8])

$$
\boxed{\Phi_{\xi\xi} = \Phi_\xi\Phi_{\eta\eta} - \Phi_\eta\Phi_{\xi\eta} - \alpha}
$$

by $\Phi \mapsto \Phi - \alpha\xi^2/2$.

6. Summary of results

Below, a concise exposition of the obtained results is given[2] in Table 6 on p. 19.

Acknowledgements

The authors are grateful to E. Ferapontov, M. Marvan, and A. Sergyeyev for discussions. Computations of symmetry algebras were fulfiled using the Jets sofware, [9].

[2]We use the notation $\infty^{k\cdot\tau}$ to indicate the infinite-dimensional component corresponding to k arbitrary functions in τ

Eqn	$\dim(\operatorname{sym}\mathscr{E})$	Reductions	Comments
(10)	$1+\infty^{2\cdot x}+\infty^{2\cdot z}$	$X\Phi_{xz}-X'\Phi_z=0$	Solves explicitly
		$2\Phi=\Phi\Phi_{xz}-\Phi_x\Phi_z$	Transforms to the Liouville equation
		$\Phi_{\xi\xi}=X'\Phi_\xi-X\Phi_{x\xi}$	Solves explicitly
		$\Phi_{\xi\xi}=\Phi_x\Phi_\xi-\Phi\Phi_{x\xi}$	
		$(1+Z\Phi_z)\Phi_{\xi\xi}=Z\Phi_\xi\Phi_{\xi z}+Z'\Phi_\xi^2$	Solves explicitly
		$(1+\xi\Phi_z)\Phi_{\xi\xi}-\xi\Phi_\xi\Phi_{\xi z}+\Phi_\xi\Phi_z=0$	
		$\Phi_{\xi\xi}=\Phi_\xi\Phi_{\eta\eta}-\Phi_\eta\Phi_{\xi\eta}$	Linearizes by the Legendre transformation
		$\Phi_\eta\Phi_{\xi\eta}-\Phi_\xi\Phi_{\eta\eta}=e^\eta\Phi_{\xi\xi}$	
(20)	$1+\infty^{1\cdot y}+\infty^{3\cdot t}$	$\Phi_{yt}=T\Phi_y$	Solves explicitly
		$\Phi_{yt}=2\Phi\Phi_y$	Reduces to the Riccati equation
		$(\alpha\xi+\Phi_\xi)\Phi_{\xi y}-\Phi_y(\Phi_{\xi\xi}+2\alpha)=0$	Solves explicitly for $\alpha=0$
		$T\Phi_{xx}=T'$	Solves explicitly
		$\left(\frac{T'}{T}\xi+\bar{T}\right)\Phi_{\xi\xi}+\Phi_{\xi t}=0$	Solves explicitly
		$\Phi_{\xi t}=4\Phi\Phi_\xi-\xi\Phi_\xi^2+2\xi\Phi\Phi_{\xi\xi}$	
		$\Phi_{\eta\eta}+(\alpha\xi+\Phi_\eta)\Phi_{\xi\eta}=\Phi_\eta(2\alpha+\Phi_{\xi\xi})$	Linearizes by the for $\alpha=0$
(27)	$\infty^{2\cdot x}+\infty^{1\cdot y}+\infty^{1\cdot t}$	$\Phi_{yt}=0$	Solves explicitly
		$\bar{X}\Phi_{xt}-\bar{X}'\Phi_t=0$	Solves explicitly
		$\bar{X}\Phi_{xy}-\bar{X}'\Phi_y=0$	Solves explicitly
		$\Phi_{\xi\xi}=\bar{X}\Phi_{x\xi}-\bar{X}'\Phi_\xi$	Solves explicitly
		$(1+\Phi_\xi)\Phi_{y\xi}=\Phi_y\Phi_{\xi\xi}$	Solves explicitly
		$(1+\Phi_\xi)\Phi_{t\xi}=\Phi_t\Phi_{\xi\xi}$	Solves explicitly
		$\Phi_\eta\Phi_{\xi\xi}+(\Phi_\eta-\Phi_\xi-1)\Phi_{\xi\eta}-\Phi_\xi\Phi_{\eta\eta}=0$	Linearizes by the Legendre transformation
(31)	$2+\infty^{4\cdot t}$	$\Phi_{yy}=(T'-T^2)y+T\bar{T}-\bar{T}'$	Solves explicitly
		$\Phi_{yy}=\frac{2\Phi_y-T'}{y+T}+T''+\frac{\bar{T}^2}{(y+T)^3}$	Solves explicitly
		$\left(\left(\frac{T'}{T}+2\alpha T\right)\xi+\bar{T}^2+\bar{\bar{T}}\right)\Phi_{\xi\xi}$	Solves explicitly
		$-\Phi_{\xi t}-\alpha T\Phi_\xi+\bar{T}'+2\alpha T\bar{T}=0$	
		$(4\xi^2-3\Phi)\Phi_{\xi\xi}-\Phi_{\xi t}-6\xi\Phi_\xi+\Phi_\xi^2+6\Phi=0$	
		$\Phi_{\xi\xi}=(\beta\xi-\Phi_\eta)\Phi_{\xi\eta}+(2\beta\eta+\Phi_\xi)\Phi_{\eta\eta}-\beta\Phi_\eta=0$	Linearizes by the Legendre transformation for $\beta=0$
		$\Phi_{\xi\xi}=(\alpha\xi+\Phi_\xi)\Phi_{\eta\eta}-\Phi_\eta\Phi_{\xi\eta}-2\alpha$	Reduces to the Gibbons-Tsarev equation, for $\alpha\neq0$
			Linearizes by the Legendre transformation for $\alpha=0$

TABLE 6. Summary of reductions

REFERENCES

[1] Adler V.E. Shabat, A.B., *Model equation of the theory of solitons*, Theor. Math. Phys., **153**, (2007) 1, 1373–1387.

[2] Błaszak, M. *Classical R-matrices on Poisson algebras and related dispersionless systems*, Phys. Lett. **A 297** (2002), 191–195.

[3] Calogero, F., Degasperis, A. *Spectral Transform and Solitons: Tools to Solve and Investigate Nonlinear Evolution Equations*. New York: North-Holland, p. 60, 1982.

[4] Dunajski M., *A class of Einstein-Weil spaces associated to an integrable system of hydrodynamic type*, J. Geom. Phys., **51** (2004), 126 –137.

[5] Dunajski M., Kryński W., *Einstein-Weyl geometry, dispersionless Hirota equation and Veronese webs*, Mathematical Proceedings of the Cambridge Philosophical Society, **157** (2014), 1, 139–150. DOI: http://dx.doi.org/10.1017/S0305004114000164 (arXiv:1301.0621).

[6] Ferapontov E.V., Khusnutdinova K.R. *Hydrodynamic reductions of multi-dimensional dispersionless PDEs: the test for integrability*, J. Math. Phys. **45** (2004), 2365. DOI: http://dx.doi.org/10.1063/1.1738951 (arXiv:nlin/0312015).

[7] Ferapontov E.V., Moss J., *Linearly degenerate PDEs and quadratic line complexes*, arXiv:1204.2777.

[8] Gibbons J., Tsarev S.P., *Reductions of the Benney equations*, Phys. Lett. **A 211** (1996) 19–24.

[9] Baran H., Marvan M., *Jets. A software for differential calculus on jet spaces and diffeties*. `http://jets.math.slu.cz`.
[10] Krasil′shchik, I.S., Lychagin, V.V., Vinogradov, A.M.: *Geometry of Jet Spaces and Nonlinear Differential Equations*, Adv. Stud. Contemp. Math. 1, Gordon and Breach, New York, London, 1986.
[11] Martínez Alonso L., Shabat A.B., *Hydrodynamic reductions and solutions of a universal hierarchy*, Theor. Math. Phys. **104** (2004), 1073–1085 (`arXiv:nlin/0312043`).
[12] M. Marvan, Private communication.
[13] Ovsienko V., *Bi-Hamiltionian nature of the equation* $u_{tx} = u_{xy}u_y - u_{yy}u_x$. Pure Appl. Math., **1** (2010), 7–17
[14] Zakharevich I., *Nonlinear wave equation, nonlinear Riemann problem, and the twistor transform of Veronese webs* (`arXiv:math-ph/0006001`).
[15] Pavlov M.V. *Integrable hydrodynamic chains*, J. Math. Phys., **44** (2003), 4134–4156.
[16] Pavlov M.V. *The Kupershmidt hydrodynamics chains and lattices*, Intern. Math. Research Notes, **2006** (2006), article ID 46987, 1–43.

Mathematical Institute, Silesian University in Opava, Na Rybníčku 1, 746 01 Opava, Czech Republic
E-mail address: `Hynek.Baran@math.slu.cz`

Independent University of Moscow, B. Vlasevsky 11, 119002 Moscow, Russia & Mathematical Institute, Silesian University in Opava, Na Rybníčku 1, 746 01 Opava, Czech Republic
E-mail address: `josephkra@gmail.com`

Faculty of Applied Mathematics, AGH University of Science and Technology, Al. Mickiewicza 30, Kraków 30-059, Poland
E-mail address: `morozov@agh.edu.pl`

Mathematical Institute, Silesian University in Opava, Na Rybníčku 1, 746 01 Opava, Czech Republic
E-mail address: `Petr.Vojcak@math.slu.cz`